



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

When  $n = 1$ ,  $U_1 = 1$ , and when  $n = 2$ ,  $U_2 = 4$ ; then (5) gives for determining  $C_1$  and  $C_2$ ,

$$C_1(2 + \sqrt{3}) + C_2(2 - \sqrt{3}) = 1,$$

$$C_1(2 + \sqrt{3})^2 + C_2(2 - \sqrt{3})^2 = 4,$$

giving  $C_1 = \frac{1}{6}\sqrt{3}$  and  $C_2 = -\frac{1}{6}\sqrt{3}$ .

(1) then becomes  $U_n = \frac{1}{6}\sqrt{3}\{(2 + \sqrt{3})^n - (2 - \sqrt{3})^n\}$ ,

the general term. The usual theory for the sum of  $n$  terms gives

$$S_n = C + C_1 m_1 \frac{m_1^n}{m_1 - 1} + C_2 m_2 \frac{m_2^n}{m_2 - 1}.$$

Substituting the values of  $m_1$ ,  $m_2$ ,  $C_1$ ,  $C_2$ , we have

$$(2) \quad S_n = C + \frac{1}{6}\sqrt{3} \left\{ \frac{(2 + \sqrt{3})^{n+1}}{\sqrt{3} + 1} + \frac{(2 - \sqrt{3})^{n+1}}{\sqrt{3} - 1} \right\}.$$

When  $n = 1$ ,  $S_n = 1$ , and (2) gives  $C = -\frac{1}{2}$ , and this in (2) gives the required sum.

Also solved by AMELIA BENSON, G. W. HARTWELL, E. B. ESCOTT, A. M. HARDING, N. P. PANDYA, and O. S. ADAMS.

## GEOMETRY.

### 489. Proposed by NATHAN ALTSHILLER, The University of Oklahoma.

The parallels to the asymptotes  $a$ ,  $b$  of a given hyperbola, drawn from a variable point of the curve, meet  $a$  and  $b$  in  $P$ ,  $Q$  respectively. The line  $PQ$  envelops an hyperbola whose asymptotes are  $a$  and  $b$ .

### I. SOLUTION BY E. J. OGLESBY, Williamsburg, Virginia.

Take the asymptotes  $a$ ,  $b$  as the axes of coördinates. Then the equation of the hyperbola may be taken as  $xy = c^2$  and the coördinates of the variable point on the hyperbola as  $(ct, c/t)$  in terms of the parameter  $t$ .  $P$  is the point  $(ct, 0)$ , and  $Q$  is  $(0, c/t)$ .

The equation of  $PQ$  may be written

$$(1) \quad t^2y - ct + x = 0.$$

We find the envelope of (1) by applying the condition that this equation shall have equal roots in the parameter  $t$ .

Hence, we have

$$(-c)^2 - 4yx = 0, \quad \text{or} \quad xy = c^2/4,$$

which is an hyperbola having  $a$  and  $b$  as asymptotes.

### II. SOLUTION BY THE PROPOSER.

*The tangents  $a$ ,  $b$  to a given conic at the points  $A$ ,  $B$ , are met by the lines  $BM$ ,  $AM$  joining  $A$  and  $B$  to a variable point  $M$  of the curve, in the points  $P$ ,  $Q$  respectively. The line  $PQ$  envelops a conic having a double contact with the given curve at the points  $A$ ,  $B$ .*

Indeed, the lines  $AM$ ,  $BM$  describe two projective pencils, hence their sections by the lines  $a$ ,  $b$  are two projective ranges.

$$(P\cdots) \asymp B(M\cdots) \asymp A(M\cdots) \asymp (Q\cdots).$$

Consequently the line  $PQ$  envelops a conic tangent to  $a$  and  $b$ . To the point  $(ab)$  considered as an element of  $a$  and  $b$  in turn correspond, in the ranges  $(Q\cdots)$  and  $(P\cdots)$ , the points  $B$  and  $A$ , these points are therefore the points of contact of  $a$  and  $b$  with the envelope, which proves the proposition.

If for  $A$ ,  $B$  are taken some remarkable points of the conic, special cases of this general proposition are obtained. For example, if  $A$  be the point at infinity of a parabola, and  $B$  its vertex, the

proposition takes the following form: *The diameter passing through a variable point of a parabola, meets the tangent at the vertex in the point P. The parallel through P to the line joining M to the vertex of the parabola, envelops another parabola having the same vertex and the same axis as the given curve.*

The proposed problem is another special case of this general proposition, namely when both A and B are at infinity.

The duals of the three propositions are, in order:

*The points of intersection of two fixed tangents to a given conic, with a variable tangent to the same curve, are projected from the points where the fixed tangents touch the conic. The point of intersection of the two projecting lines describes a conic having a double contact with the given curve.*

*From a variable point of the tangent at the vertex of a given parabola, are drawn the diameter and the tangent to the curve. The point of intersection of the diameter with the parallel to the tangent through the vertex of the curve, describes a parabola having the same axis and the same vertex as the given curve.*

*The parallels to the asymptotes of a given hyperbola drawn through the points of intersection of the latter lines with a variable tangent to the curve, intersect in a point whose locus is an hyperbola having the same asymptotes as the given curve.*

Also solved by O. S. ADAMS, CLARA L. BACON, J. W. CLAWSON, A. M. HARDING, HORACE OLSON, PAUL CAPRON, G. W. HARTWELL, R. M. MATHEWS, and N. P. PANDYA.


**490. Proposed by ELMER E. MOOTS, University of Arizona.**

In any quadrilateral  $ABCD$ , let  $AC$  and  $BD$  be the diagonals intersecting at  $K$ . On  $AC$ , lay off  $CR$  equal to  $AK$ . Join  $B$  and  $R$ . Connect the middle point  $G$  of  $BR$  with  $D$ . On  $GD$  lay off  $GM$  equal to  $\frac{1}{3}GD$ . Show that  $M$  is the center of gravity of the quadrilateral.

SOLUTION BY A. M. HARDING, University of Arkansas.

It is evident that  $M$  is the center of gravity of the triangle  $BDR$ . Hence it will be sufficient to prove that the triangle  $BDR$  and the quadrilateral  $ABCD$  have the same center of gravity.

Let  $O$  be the mid-point of  $RK$ , then it will also be the mid-point of  $CA$ . Then  $G_1$  is the center of gravity of the triangles  $RDK$  and  $CDA$ , and  $G_2$  is the center of gravity of the triangles  $RBK$  and  $CBA$  where  $OG_1 = \frac{1}{3}OD$  and  $OG_2 = \frac{1}{3}OB$ .



Since

$$\frac{\triangle RDK}{\triangle RBK} = \frac{\triangle CDA}{\triangle CBA},$$

it follows that the center of gravity of the triangle  $BDR$  will also be the center of gravity of the quadrilateral  $ABCD$ .

Also solved by J. W. CLAWSON, O. S. ADAMS, and N. P. PANDYA.

**491. Proposed by N. P. PANDYA, Sojitra, India.**

In a triangle  $mx = b$  and  $nx = c$ , determine a relation between  $m$ ,  $n$ ,  $x$ ,  $A$  and  $s$ , and solve it for  $x$ .